

Polymerization Studies of Sickle Cell Hemoglobin Incubated in Aqueous Leaf Extract of *Nicotiana tabacum* Product

By

P.C. Chikezie, A.R. Akuwudike and C.M. Chikezie

ISSN 0970-4973 (Print)

ISSN 2319-3077 (Online/Electronic)


Volume 30
No.1 (2013)

J. Biol. Chem. Research
Volume 30 (1) 2013 Pages No. 158-168

**Journal of
Biological and
Chemical Research**

(An International Journal of Life Sciences and Chemistry)

Published by Society for Advancement of Sciences®





JBCR

[http:// www.jbcr.in](http://www.jbcr.in)

jbicrchemres@gmail.com info@jbcr.in

RESEARCH PAPER

Received: 21/12/2012 Revised: 18/02/2013 Accepted: 20/02/2013

Polymerization Studies of Sickle Cell Hemoglobin Incubated in Aqueous Leaf Extract of *Nicotiana tabacum* Product

P.C. Chikezie, A.R. Akuwudike and C.M. Chikezie

Department of Biochemistry, Imo State University, Owerri, Nigeria

ABSTRACT

The present study seeks to ascertain the capacity of aqueous extract of N. tabacum to alter and interfere with polymerization of deoxygenated sickle cell haemoglobin (deoxyHbS) molecules in vitro. Spectrophotometric method was used to measure level of sodium metabisulphite induced polymerization of deoxyHbS molecule incubated in aqueous extract of N. tabacum for 180 s. The polymerization profile of deoxyHbS molecules of control and test samples showed increasing level of polymerization with progression of experimental time. At experimental time $t > 30$ s, level of polymerization of control sample ranged between $60.9 \pm 0.76 - 100 \pm 1.05\%$, whereas, the test samples ranged in the following corresponding order: $[N. tabacum] = 0.8 \text{ mg}/100 \text{ mL}$, $65.6 \pm 0.93 - 176.0 \pm 4.26\%$; $[N. tabacum] = 1.0 \text{ mg}/100 \text{ mL}$, $75.7 \pm 1.07 - 192.9 \pm 5.03\%$ and $[N. tabacum] = 2.0 \text{ mg}/100 \text{ mL}$, $135.9 \pm 5.04 - 297.2 \pm 19.14\%$. Activation of deoxyHbS polymerization by aqueous extract of N. tabacum increased in a concentration dependent manner and duration of incubation. Specifically, at $t = 180$ s, $[N. tabacum] = 2.0 \text{ mg}/100 \text{ mL}$ caused highest level of activation of deoxyHbS polymerization ($197.2 \pm 19.14\%$); an increase in %polymerization in a ratio of 1: 4.5 (approx.) compared with $[N. tabacum] = 0.8 \text{ mg}/100 \text{ mL}$ at $t = 30$ s. The study showed that aqueous extract of N. tabacum exacerbated polymerization of deoxyHbS molecules in a concentration and time dependent manner. Therefore, N. tabacum in the present experimental form did not exhibit therapeutic potentials for management and alleviation of sickling disorder.

Keywords: Polymerization, Sickle cell haemoglobin, N. tabacum, Sodium metabisulphite.

INTRODUCTION

The sickle cell gene (β^S) occurs widely throughout Africa, part of Asia, the Arabian Peninsula and part of Southern Europe. In Africa, there are two areas with very high frequencies of β^S . One includes Nigeria, Ghana, and the other, Gabon and Zaire (Waniscoat, 1987; Uzoegwu and Onwurah, 2003). Sickle cell anaemia (SCA) occurs with a much lower incidence in part of Italy, Greece, the Middle East and India. The homozygous state of SCA is associated with complications and reduced life expectancy (Kutlar, 2007; Frenette and Atweh, 2007).

The sickle cell haemoglobin (HbS) variant is caused by a point mutation affecting the coding sequence of the β -globin gene resulting in a substitution of glutamic acid by valine at the sixth position of β -globin chains (Kutlar, 2007; Bianchi *et al.*, 2009). Under low oxygen tension, deoxyHbS molecules polymerize into microfibrils parallel to each other with concomitant reduced solubility (forms gel) and resultant erythrocyte membrane deformity and damage (Xue *et al.*, 2002). Therapeutic approaches for the treatment and management of SCA disease include induction of fetal hemoglobin expression (Bianchi *et al.*, 2009; Fathallah and Atweh, 2006), bone marrow transplantation (Billing, 1990; Walters *et al.*, 2000), and the use of pharmacological agents that interact either non-covalently or covalently with HbS molecules (Xue *et al.*, 1999; Abdulmalik *et al.*, 2005) to retard or inhibit haemoglobin aggregation and polymerization.

The tobacco plant (*Nicotiana tabacum*) has been used for several years and purposes, irrespective of the location of human races. Cigarettes, in spite of the health risk associated with its usage constitute the largest share of manufactured tobacco products in the world, accounting for 96% of total sales. (Adeniyi *et al.*, 2010). The chemical compositions of *N. tabacum* leaf extract have been previously reported (Andersen and Kasperbauer, 1973; Siddiqui *et al.*, 2004; Chitra and Sivaranjani, 2012). The prospects of utilization of *N. tabacum* products for purposes that do not pose health risk are now widely encouraged and advertised (Mugisha-Kamatenesi *et al.*, 2008). There are reports and claims by several authors on the medicinal benefits (Chitra and Sivaranjani, 2012; Nacoulma *et al.*, 2012) and biologic activities, such as ovicidal, repellent and insecticidal activities against various insect species (Isman, 2006; Xue *et al.*, 2009; Farsani *et al.*, 2011) by leaf extracts of *N. tabacum*. Application of herbal remedies for the management of SCA has been widely reported (Ekeke and Shode, 1985; Kade *et al.*, 2003; Okpuzor *et al.*, 2008; Imaga *et al.*, 2010). *In vitro* studies have revealed that plant extracts altered the polymerization of deoxyHbS molecules (Imaga *et al.*, 2010; Chikezie, 2011). Therefore, the present study seeks to ascertain the capacity of aqueous extract of *N. tabacum* to alter and interfere with the polymerization of deoxyHbS molecules *in vitro*.

MATERIAL AND METHODS

Collection and preparation of *N. tabacum* leaf extract

Preparation of aqueous extract of *N. tabacum* was according to the methods earlier described by Chikezie and Uwakwe, (2011). Twenty five grams (25 g) of processed leaves of *N. tabacum* obtained from the popular cigarette brand Benson & Hedges™ were collected in desiccators and allowed to dry for 72 h to become crispy. The dried specimen was ground in ceramic mortar and pestle into fine powder. The pulverized specimen was suspended in 100 mL of distilled water and allowed to stand for 6 h at 37°C. Aqueous extract of *N. abacus* was obtained by filtration with Whatman No. 2 filter paper. Finally, the extract was concentrated in a rotary evaporator at 50°C and dried in vacuum desiccators. The yield was calculated to be 5.14% (1.29 g; w/w). The extract was finally suspended in 50 mL phosphate buffered saline (PBS) solution osmotically equivalent to 0.9 g/100 mL NaCl {NaCl (9.0 g), Na₂HPO₄·2H₂O (1.71 g) and NaH₂PO₄·2H₂O (0.243 g)/100 mL; pH =7.4} (Ibegbulem and Chikezie, 2012). The extract was kept at 4°C in a refrigerator for at least 24 h before subsequent tests. Concentration equivalents of 0.8, 1.0 and 2.0 mg/100 mL aqueous extracts of *N. abacus* were used for polymerization studies.

Ethics: The institutional review board of the Department of Biochemistry, Imo State university, Owerri, Nigeria, granted approval for this study and all participants involved signed an informed consent form. This study was in accordance with the ethical principles that have their origins in the Declaration of Helsinki. Individuals drawn were from Imo State University, Owerri and environs. The research protocols were in collaboration with registered and specialized clinics and medical laboratories.

Collection of Blood Samples/Preparation of Erythrocyte Haemolysate

Five milliliters (5.0 mL) of venous blood obtained from the volunteers by venipuncture was stored in EDTA anticoagulant tubes. Blood samples were from patients attending clinics at the Federal Medical Center (FMC), Imo State University Teaching Hospital (IMSUTH), Orlu, St. John Clinic/Medical Diagnostic Laboratories, Avigram Medical Diagnostic Laboratories, and Qualitech Medical Diagnostic Laboratories. These centers are located in Owerri, Imo State, Nigeria.

The erythrocytes were separated from plasma and washed by centrifugation method as described by Tsakiris *et al.*, (2006) with minor modification according to Penningset *et al.*, (1999). Blood volume of 4.0 mL was introduced into centrifuge test tubes containing 4.0 mL of buffer solution pH = 7.4: 250 mM Tris (hydroxyl methyl) amino ethane-HCl (Tris-HCl)/ 140 mM NaCl/ 1.0 mM MgCl₂/ 10 mM glucose and centrifuged at 4000 *rpm* for 10 min using a centrifuge (B. Bran Scientific and Instrument Company, England). The plasma and buffy coat (supernatant) were carefully removed with Pasteur pipette. This process was repeated until the supernatant became clear. The pelleted erythrocytes were suspended in PBS (pH=7.4) to obtain approximately 10% haematocrit and stored at 4°C for 24 h.

The washed erythrocytes were lysed by freezing/thawing as described by Galbraith and Watts, (1980) and Kamberet *et al.*, 1984. The erythrocyte haemolysate was used for polymerization studies.

Polymerization Studies

Sodium metabisulfite {Na₂S₂O₅; (BDH, UK)}-induced polymerization of HbS molecules was ascertained as described previously by Iwu et al., (1988) with minor modification according to Chikezie *et al.*, (2010). The underlying principle is that HbS molecules undergo gelation when deprived of oxygen, transiting to deoxyHbS molecules; Na₂S₂O₃ was used as the reductant. The level of polymerization was measured by recording increasing absorbance of the assay mixture with progression of time. A 0.1 mL of HbS haemolysate was introduced into a test tube, followed by 0.5 mL of PBS and 1.0 mL of distilled water. The mixture was transferred into a cuvette and 3.4 mL of 2 g/100 mL aqueous solution of Na₂S₂O₅ was added. The absorbance ($\lambda_{\text{max}} = 700 \text{ nm}$) of the assay mixture was measured with a spectrophotometer (SPECTRONIC 20, Labtech—Digital Blood Analyzer®) at every 30 s for 180 s (control assay). This procedure was repeated by substituting distilled water with 1.0 mL of corresponding three increasing concentrations (0.8, 1.0 and 2.0 mg/100 mL) of *N. tabacum* aqueous extracts (test assay).

Calculations

Percentage polymerization was calculated according to Chikezie *et al.* (2010), thus;

$$\% \text{Polymerization} = \frac{A_{t/c} \cdot 100}{A_{c180th s}}$$

Where

$A_{t/c}$ = Absorbance of test/control assay at time = t s.

$A_{c180th s}$ = Absorbance of control assay at the 180th s.

Cummulative polymerization (%polymerization.second) was evaluated using the Simpson's Rule. Thus:

$$f(X_1)h_1 + f(X_2)h_2 + \dots + f(X_n)h_n$$

Area under the curve (AUC) of the plot of %polymerization versus time (s) is given by:

$$AUC(\% \text{polymerization. second}) = \frac{h}{2} (y_n + 2y_{n-1} + 2y_{n-2} + 2y_{n-3} + \dots) \text{Equation 1}$$

Where

h = time intervals (30 s).

y = %polymerization at corresponding time interval.

$$\text{Thus: } AUC(\% \text{polymerization. second}) = \frac{h}{2} (y_6 + 2y_5 + 2y_4 + 2y_3 + 2y_2 + y_1) \text{Equation 2}$$

RESULTS

The results illustrated in Figure 1 showed that deoxyHbS molecules of the control and test samples showed increasing level of polymerization with progression of experimental time.

At experimental time $t > 30$ s, level of polymerization of control sample ranged between $60.9 \pm 0.76 - 100 \pm 1.05\%$, whereas, the test samples ranged in the following corresponding order: $[N. tabacum] = 0.8 \text{ mg}/100 \text{ mL}$, $65.6 \pm 0.93 - 176.0 \pm 4.26\%$; $[N. tabacum] = 1.0 \text{ mg}/100 \text{ mL}$, $75.7 \pm 1.07 - 192.9 \pm 5.03\%$ and $[N. tabacum] = 2.0 \text{ mg}/100 \text{ mL}$, $135.9 \pm 5.04 - 297.2 \pm 19.14\%$.




Figure 1. Levels of DeoxyHbS polymerization of control and in presence of *N. tabacum* with time.

Therefore, the three experimental concentrations of *N. tabacum* caused activation of deoxyHbS polymerization in the order: $[N. tabacum] = 0.8 \text{ mg/100 mL} < 1.0 \text{ mg/100 mL} < 2.0 \text{ mg/100 mL}$. Furthermore, cumulative level of deoxyHbS polymerization was in the order: Control < $[N. tabacum] = 0.8 \text{ mg/100 mL} < 1.0 \text{ mg/100 mL} < 2.0 \text{ mg/100 mL}$ (Table 1).

Table 1. Cumulative level of polymerization of deoxyHbS within experimental $t = 180 \text{ s}$.

$[N. tabacum]$ mg/100 mL	Control	0.8	1.0	2.0
AUC (%polymerization.second) $\times 10^3$	13.2	20.0	21.3	52.0

Table 2 showed that activation of deoxyHbS polymerization by aqueous extract of *N. tabacum* increased in a concentration dependent manner and duration of incubation. Specifically, at $t = 180 \text{ s}$, $[N. tabacum] = 2.0 \text{ mg/100 mL}$ caused highest level of activation of deoxyHbS polymerization ($197.2 \pm 19.14\%$). The value represented an increase of %polymerization in a ratio of 1: 4.5 (approx.) compared with $[N. tabacum] = 0.8 \text{ mg/100 mL}$ at $t = 30 \text{ s}$.

Table 2. Activation of DeoxyHbS polymerization in the presence aqueous extracts of *N. tabacum* within intervals of $t = 30 \text{ s}$.

Percentage activation of deoxyHbS polymerization						
Time (s)/ $[N. tabacum]$ mg/100 mL	30	60	90	120	150	180
0.8	4.7 $\pm$ 0.93	17.5 $\pm$ 1.64	60.3 $\pm$ 2.15	50.5 $\pm$ 2.03	60.7 $\pm$ 3.14	76.0 $\pm$ 4.26
1.0	14.8 $\pm$ 1.07	30.6 $\pm$ 2.08	49.5 $\pm$ 3.15	65.3 $\pm$ 4.04	86.6 $\pm$ 4.16	92.9 $\pm$ 5.03
2.0	75.0 $\pm$ 5.04	97.6 $\pm$ 9.04	133.0 $\pm$ 10.0	150.1 $\pm$ 17.0	167.1 $\pm$ 19.0	197.2 $\pm$ 19.14
			4	4	1	

The results are means ($\bar{X}$) $\pm$ SD of six ($n = 6$) determinations.

When $t > 30 \text{ s}$, deoxyHbS incubated in aqueous extract $[N. tabacum] = 2.0 \text{ mg/100 mL}$ exhibited %polymerization > 100%. Likewise, %polymerization > 100% in the presence of 0.8 and 1.0 mg//100 mL of *N. tabacum* occurred when $t > 60 \text{ s}$ and $t > 30 \text{ s}$ respectively.

DISCUSSION

The nature and mode of interaction between biochemical agents and contact points of deoxyHbS molecules, in addition to erythrocyte membrane structural components are crucial determining factors in the capability of these agents to retard/inhibit polymerization of deoxyHbS molecules, to alleviate and prevent erythrocyte sickling disorders (Oyewole et al., 2008; Chikezie, 2011; Chikezie and Uwakwe, 2011; Ibegbulemet et al., 2011). Contrary to reports by previous authors and researchers on the anti-polymerization (Imagaet et al., 2010; Chikezie, 2011) and claims of anti-sickling potencies (Ekeke and Shode, 1985; Kadeet et al., 2003; Okpuzoret et al., 2008; Imagaet et al., 2010; Ibegbulemet et al., 2011) of plant extracts, the present report showed that aqueous extract of *N. tabacum* exacerbated polymerization of deoxyHbS molecules. Chitra and Sivaranjani, (2012) have previously reported that processed *N. tabacum* (leaf powder) was devoid of flavonoids and terpenoids. These two phytochemicals have been implicated in inhibiting polymerization of deoxyHbS molecules and by extension exhibited anti-sickling activity on human HbSS erythrocytes *in vitro* (Fall et al., 1999; Ibegbulemet et al., 2011). However, it is worthwhile to note that the biologic action of *N. tabacum* on HbS molecules reported here and the capacity plant extracts to alter erythrocytes physiochemical properties is an additive effect of combination of the plant components (Mikstacka et al., 2010).

Haemoglobin polymerization and subsequent sickling of erythrocytes is favoured when HbS molecules are deoxygenated (Martin, 1983; Poillon et al., 1998). In the present *in vitro* study, the aggravation of polymerization of deoxyHbS molecules by the three experimental concentrations of *N. tabacum* was an obvious reflection of the capability of the plant extract to act in synergy with the hypoxemic agent ($\text{Na}_2\text{S}_2\text{O}_5$), thereby facilitating more rapid rate of polymerization of HbS molecules compared with the control assay (Figure 1). Conversely, chemical compounds and phytochemicals that serve to stabilize the R-state of HbS molecules with resulting increase oxygen/haemoglobin affinity will cause reduction in the rate and degree of HbS polymerization (Oyewole et al., 2008; Chikezie, 2011). Aqueous extracts of *N. tabacum* in the present experimental form did not demonstrate R-state stabilizing property *in vitro* and therefore, did not exhibit therapeutic potentials for the management and alleviation of sickling disorder.

ACKNOWLEDGEMENT

We thank Professor A. A. Awake, Professor of Medical Biochemistry, University of Port Harcourt, for providing us with useful information for the preparation of this article. We also appreciate the efforts of our undergraduate students.

REFERENCES

- Abdulmalik, O., Safo, M. K., Chen, Q., Yang, J., Brugnara, C., et al., 2005. 5-hydroxymethyl-2-furfural modifies intracellular sickle haemoglobin and inhibits sickling red blood cells. *British Journal of Haematology*. 128(4), 552-561.
- Adeniyi, P. A. O., Ghazal, O. K., Musa, A. A. and Caxton-Martins, E. A. 2010. The neurobehavioural effects of smoke and ethanolic extract of *Nicotiana tabacum* leaves exposure in mice. *Research Journal of Animal Sciences*. 4(4), 99-102.
- Andersen, G. and Kasperbauer, M. J. 1973. Chemical composition of tobacco leaves altered by near-ultraviolet and intensity of visible light. *Plant Physiology*. 51, 723-726.
- Bianchi, N., Zuccato, C., Lampronti, I., Borgatti, M. and Gambari, R. 2009. Fetal hemoglobin inducers from the natural world: A novel approach for identification of drugs for the treatment of β -thalassemia and sickle-cell anemia. *eCAM*. 6(2), 141-151; doi:10.1093/ecam/nem139.
- Billings, F. T. 1990. Treatment of sickle cell anemia with bone marrow transplantation-pros and cons. *Transaction of American Clinical Climatology Association*. 101, 8-20.
- Chikezie, P. C. 2011. Sodium metabisulphite induced polymerization of sickle cell haemoglobin (HbS) incubated in extracts of three medicinal plants (*Anacardium occidentale*, *Psidium guajava* and *Terminalia catappa*). *Pharmacognosy Magazine*. 7(26), 126-132.
- Chikezie, P. C. and Uwakwe A. A. 2011. Membrane stability of sickle erythrocytes incubated in three medicinal plants: *Anacardium occidentale*, *Psidium guajava* and *Terminalia catappa*. *Pharmacognosy Magazine*. 7(26), 121-125.
- Chikezie, P. C., Chikezie, C. M. and Amaragbulem, P. I. 2010. Effect of antimalarial drugs on polymerization of sickle cell hemoglobin (HbS). *Turkish Journal of Biochemistry*. 35(1), 41-44.
- Chitra, S. and Sivaranjani, K. 2012. A comparative phytochemical analysis of tobacco and its natural extract- an eccentric approach. *International Journal of Pharmacy and Pharmaceutical Sciences*. 4(2), 1-2.
- Ekeke, G. I. and Shode, F. O. 1985. The reversion of sickled cells by *Cajanus cajan*. *Planta Medica*. 6, 504-507.
- Fall, A. B., Vanhaelen-Fastre, R., Vanhaelen, M., Lo, I., Toppet, M., Ferster, A. and Fondu, P. 1999. *In vitro* antisickling activity of a rearranged limonoid isolated from *Khaya senegalensis*. *Planta Medica*. 65(3), 209-212.
- Farsani, N. S., Zamani, A. A., Abbasi, S. and Kheradmand, K. 2011. Insecticidal effects of two plant aqueous extracts against second instar larvae of *Lycoriella auripila* (Diptera: Sciaridae). *World Academy of Science, Engineering and Technology*. 58, 439-441.
- Fathallah, H. and Atweh, G. F. 2006. DNA hypomethylation therapy for hemoglobin disorders: molecular mechanisms and clinical applications. *Blood Review*. 20, 227-234.

- Frenette, P. S. and Atweh, G. F. 2007. Sick cell disease: old discoveries, new concepts, and future promise. *Journal of Clinical Investigation*.117, 850–858.
- Galbraith, D. A. and Watts, D. C. 1980. Changes in some cytoplasmic enzymes from red cells fractionated into age groups by centrifugation in Ficoll™/Trisil™ gradients: Comparison of normal human and patients with Duchenne muscular dystrophy. *Biochemistry Journal*.191, 63-70.
- Ibegbulem, C. O. and Chikezie, P. C. 2012. Serum lipid profile of rats (*Rattus norvegicus*) fed with palm oil and palm kernel oil-containing diets. *Asian Journal of Biochemistry*.7(1), 46-53.
- Ibegbulem, C. O., Eyong, E. U. and Essien, E. U. 2011. Polymerization inhibition activity of *Raphia hookeri* palm sap and its effect on osmotic fragility of sickle cell red blood cells. *Journal of Medicinal Plants Research*.5(17), 4212-4217.
- Imaga, N. O. A., Gbenle, G. O., Okochi, V. I., Adenekan, S. O., Edeghon, S. O., Kehinde, M. O., Bamiro, S. B., Ajiboye, A. and Obinna, A. 2010. Antisickling and toxicological profiles of leaf and stem of *Parquetinanigrescens* L. *Journal of Medicinal Plants Research*.4(8), 639-643.
- Isman, M. B. 2006. Botanical insecticides, deterrents, and repellents in modern agriculture and increasingly regulated world. *Annual Review of Entomology*.51, 45–66.
- Iwu, M. N., Igboko, A. O., Onwubiko, H. and Ndu, U. E. 1988. Effect of *Cajanus cajan* on gelation and oxygen affinity of sickle cell hemoglobin. *Journal of Ethnopharmacology*.20, 99-104.
- Kade, I. J., Kotila, O. O., Ayeleso, A. O., Olaleye, A. A. and Olawoye, T. L. 2003. Antisickling properties of *Parquetinanigrescens*. *Biomedical Research (Allgarh)*.14, 185-188.
- Kamber, K., Poyiagi, A. and Delikonstantinos, G. 1984. Modifications in the activities of membrane-bound enzymes during *in vivo* ageing of human and rabbit erythrocytes. *Comparative Biochemistry and Physiology* B.77B: 95-99.
- Kutlar A. 2005. Sick cell disease: A multigenic perspective of a single gene disorder. *Medical Principles and Practice* 14, 15-19.
- Martin, D. W. 1983. Structure and function of a protein - haemoglobin. In: Martin, D. W., Mayes, P. A. and Rodwell, V. W. editors. *Harper's Review of Biochemistry*. 9th ed. California: Lange. Medical Publications.
- Mikstacka, R., Agnes, M. R. and Ewa, I. 2010. Antioxidant effect of trans-resveratrol, pterostilbene, quercetin and their combinations in human erythrocytes. *in vitro. Plant Foods and Human Nutrition*.65, 57–63.
- Mugisha-Kamatenesi, M., Deng, M. L., Ogendo, J. O., Omolo, E. O., Mihale, M. J., Otim, M. J. Buyungo, J. P. and Bett, P. K. 2008. Indigenous knowledge of field insect pests and their management around lake Victoria basin in Uganda. *African Journal of Environmental Science and Technology*. 2(8), 342-348.

- Nacoulma, A. P., Compaoré, M., de Lorenzi, M., Kiendrebeogo, M. and Nacoulma, O. G. 2012. *In vitro* Antioxidant and anti-inflammatory activities of extracts from *Nicotianatabacum* L. (Solanaceae) leafy galls induced by *Rhodococcus fascians*. *Journal of Phytopathology*. 160(10), doi: 10.1111/j.1439-0434.2012.01953.x
- Okpuzor, J., Adebisin, O., Ogbunugafor, H. and Amadi, I. 2008. The potential of medicinal plants in sickle cell disease control: A review. *International Journal of Biomedicine and Health Science*. 4(2), 47-55.
- Oyewole, O. I., Malomo, S. O. and Adebayo, J. O. 2008. Comparative studies on antisickling properties of thiocyanate, tellurite and hydroxyurea. *Pakistan Journal of Medical Science*. 24, 18-22.
- Pennings, H. J., Borm, P. J. A., Evelo, C. T. A. and Wouters, E. F. M. 1999. Changes in levels of catalase and glutathione in erythrocytes of patients with stable asthma, treated with beclomethasonedipropionate. *European Respiratory Journal*. 13, 1260-1266.
- Poillon, W. N., Kim, B. C. and Castro, O. 1998. Intracellular hemoglobin S polymerization and the clinical severity of sickle cell anemia. *Blood*. 91(5), 1777-1783.
- Siddiqui, M. H., Oad, F. C. and Kumar, L. 2004. Growth and chemical composition of tobacco *Nicotianatabacum* L. under different seedling height. *Asian Journal of Plant Science*. 3, 251-253.
- Tsakiris, S., Giannoulia-Karantana, A., Simintzi, I. and Schulpis, K. H. 2005. The effect of aspartame metabolites on human erythrocyte membrane acetylcholinesterase activity. *Pharmacology Research*. 53, 1-5.
- Uzoegwu, P. N. and Onwurah, A. E. 2003. Correlation of lipid peroxidation index with sickle haemoglobin concentrations in malaria-positive and negative statuses of AA, AS and SS individuals from the University of Nigeria, Nsukka community. *Research Biotechnology*. 1(1), 97-114
- Wainscoat, J. S. 1987. The origin of mutant globin genes in human population. *Acta Haematologica*. 78, 154-158.
- Walters, M. C., Storb, R. and Patience, M. 2000. Impact of bone marrow transplantation for symptomatic sickle cell disease: an interim report. *Blood*. 95, 1918-1924.
- Xu, A. S. L., Labotka, R. J. and London, R. E. 1999. Acetylation of human hemoglobin by methyl acetylphosphate: Evidence of broad regio-selectivity revealed by NMR studies. *Journal of Biological Chemistry*. 274, 26629-26632.

- Xu, X., Lockamy, V. L., Chen, K., Zhi, H., Shields, H et al., 2002. Effects of iron nitrosylation on sickle cell hemoglobin solubility. *Journal of Biological Chemistry*. 277, 36787-36792.
- Xue, M., Pang, Y., Wang, H., Li, Q. and Liu, T. 2009. Effects of four host plants on biology and food utilization of the cutworm, *Spodopteralitura*. *Journal of Insect Science*. 10(22), 1-14.

Corresponding author: Dr. P.C. Chikezie, Department of Biochemistry, Imo State University, Owerri, Nigeria.

E-mail: p_chikezie@yahoo.com **Phone:** +2348038935327.